

# MATH 223: Multivariable Calculus

## Notes on Class 2

September 14, 2022

## Analog of Straight Line In Higher Dimensions

Line:  $ax + by = c$

$$a_1x_1 + a_2x_2 = c$$

Plane:  $a_1x_1 + a_2x_2 + a_3x_3 = d$       Other Important

$$ax + by + cz = d$$


Hyperplane:  $a_1x_1 + a_2x_2 + \dots + a_nx_n = d$

Curves: CIRCLES and ELLIPSES  
and the counterparts in higher dimensions.

Recall: Graph of  $f : \mathcal{R}^1 \rightarrow \mathcal{R}^1$  is a curve in the plane.  
BUT: NOT EVERY CURVE IN THE PLANE IS THE GRAPH OF  
SUCH A FUNCTION


Vertical Line Test

## CIRCLE



Set of all points a  
fixed distance from  
a fixed point (center)  
 $\text{distance}(P, C) = r$

## ELLIPSE



Set of all points,  
sum of distances  
to pair of fixed points is constant  
 $d(P, C_1) + d(P, C_2) = r$


Distance in  $\mathcal{R}^n$

Magnitude of a vector  $\mathbf{v} = |\mathbf{v}| = \sqrt{\mathbf{v} \cdot \mathbf{v}}$

Where does this come from?

Consider  $\mathbf{v} = (v_1, v_2)$  in  $\mathcal{R}^2$  where  $\mathbf{v} \cdot \mathbf{v} = v_1^2 + v_2^2$

$\sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2}$  (Pythagorean Theorem)



Distance Between  $x$  and  $y = |x - y|$

Examine Circle in the Plane

Center  $\mathbf{C} = (a, b)$  and radius  $r$

Variable Point  $\mathbf{P} = (x, y)$

Defining Relationship:  $d(\mathbf{P}, \mathbf{C}) = r$

which means

$$|\mathbf{P} - \mathbf{C}| = r$$

$$|(x - a, y - b)| = r$$

$$\sqrt{(x - a)^2 + (y - b)^2} = r$$

$$(x - a)^2 + (y - b)^2 = r^2$$

Multiply Out:  $x^2 - 2ax + a^2 + y^2 - 2by + b^2 = r^2$

$$x^2 + y^2 - 2ax - 2by = r^2 - a^2 - b^2$$

which has the form  $x^2 + y^2 + Ax + By = C$ .

Can We Go Backwards?

Example:  $x^2 + y^2 - 6x + 16y = 71$

Complete the Squares in  $x$  and  $y$

$$(x^2 - 6x) + (y^2 + 16y) = 71$$

$$(x^2 - 6x + 9) + (y^2 + 16y + 64) = 71 + 9 + 64$$

$$(x - 3)^2 + (y + 8)^2 = 144 = 12^2$$

Circle as Image of a function  $f : \mathcal{R}^1 \rightarrow \mathcal{R}^2$

**Parametrization** with parameter  $t$

Example:  $\mathbf{f}(t) = (\cos t, \sin t), 0 \leq t \leq 2\pi$

Example:

$$\begin{cases} x = 12 \cos t + 3 \\ y = 12 \sin t - 8 \end{cases}$$

$$\begin{cases} x - 3 = 12 \cos t \\ y + 8 = 12 \sin t - 8 \end{cases}$$

$$(x - 3)^2 + (y + 8)^2 = 12^2$$

$$\mathbf{f}(t) = (12 \cos t + 3, 12 \sin t - 8)$$

## ELLIPSE

Standard Ellipse:

Center at  $(0,0)$

Foci at  $(\pm c, 0)$

Vertices  $(\pm a, 0)$  and  $(0, \pm b)$

$(a,0)$  distance from  $(c,0)$  + distance from  $(-c,0)$

$$(a - c) + (a - (-c)) = a - c + a + c = 2a$$

$(0,b)$  distance from  $(c,0)$  + distance from  $(-c,0)$

$$\sqrt{c^2 + b^2} + \sqrt{c^2 + b^2} = 2\sqrt{c^2 + b^2}$$

$$\text{Thus } 2\sqrt{c^2 + b^2} = 2a$$

$$\text{So } c^2 + b^2 = a^2 \text{ implying } c^2 = a^2 - b^2$$

$(x,y)$  distance from  $(c,0)$  + distance from  $(-c,0) = 2a$

$$\sqrt{(x - c)^2 + y^2} + \sqrt{(x + c)^2 + y^2} = 2a$$

Much algebra yields

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

Parametrization:  $\mathbf{g}(t) = (a \cos t, b \sin t), 0 \leq t \leq 2\pi$

## The Algebra

$$\sqrt{(x-c)^2 + y^2} + \sqrt{(x+c)^2 + y^2} = 2a$$

Write as  $\sqrt{(x+c)^2 + y^2} = 2a - \sqrt{(x-c)^2 + y^2}$

Square Both Sides

$$(x+c)^2 + y^2 = 4a^2 - 4a\sqrt{(x-c)^2 + y^2} + (x-c)^2 + y^2$$

Expand, Simplify, and Divide by 4

$$x^2 + 2cx + c^2 + y^2 = 4a^2 - 4a\sqrt{(x-c)^2 + y^2} + x^2 - 2cx + c^2 + y^2$$

$$4cx - 4a^2 = -4a\sqrt{(x-c)^2 + y^2}$$

$$cx - a^2 = -a\sqrt{(x-c)^2 + y^2}$$



Begin with  $cx - a^2 = -a\sqrt{(x - c)^2 + y^2}$

Square Again

$$c^2x^2 - 2a^2cx + a^4 = a^2((x - c)^2 + y^2)$$

$$c^2x^2 - 2a^2cx + a^4 = a^2x^2 - 2a^2cx + a^2c^2 + a^2y^2$$

$$c^2x^2 + a^4 = a^2x^2 + a^2c^2 + a^2y^2$$

Write as  $(a^2 - c^2)x^2 + a^2y^2 = a^4 - a^2c^2 = a^2(a^2 - c^2)$

But  $a^2 - c^2 = b^2$  so

$$b^2x^2 + a^2y^2 = a^2b^2$$

Divide by  $a^2b^2$ :

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$