


MATH 226: Differential Equations



"You know how it is! We get a warhead, *they* have to get a warhead."

CartoonStock.com



Notes on Assignment 8

Assignment Linear Algebra Computations with

Maple

Graded Project 1

Announcements

▶ Exam 1

- ▶ Wednesday
- ▶ 7 PM - ? (No Time Limit)
- ▶ Warner 101 (Here)
- ▶ No Calculators, Books, Notes, Smart Phones, etc.
- ▶ Focus on Material in Chapters 1 and 2

▶ Exam 1 Tips

- ▶ Read Overall Directions on Cover Page
- ▶ Read Each Problem Statement Carefully
- ▶ Pick "Easy" Problem To Do First
- ▶ Show All Intermediate Steps
- ▶ Include Explanations
- ▶ Pay Attention to Units
- ▶ Double Check Your Work

Today's Topic

Analysis of The Richardson Arms Race Model

Richardson Arms Race Model



Lewis F. Richardson
1881 – 1953

$x(t)$ = Arms Expenditure of Blue Nation

$y(t)$ = Arms Expenditure of Red Nation

$$x' = ay - mx + r$$

$$y' = bx - ny + s$$

where a, b, m, n are positive constants while r and s are constants.

Structure: $\vec{X}' = A\vec{X} + \vec{b}$ or $\mathbf{x}' = A\mathbf{x} + \mathbf{b}$

$$x' = ay - mx + r$$

$$y' = bx - ny + s$$

where a, b, m, n are positive constants while r and s are constants.

Find stable Lines L and L' where $x' = 0$ and $y' = 0$.

$$L : y = \frac{m}{a}x - \frac{r}{a}$$

$$L' : y = \frac{b}{n}x + \frac{s}{n}$$

Determine the Stable Point (x^*, y^*) where lines L and L' intersect.

$$ay^* - mx^* + r = 0, \quad bx^* - ny^* + s = 0$$

$$x' = ay - mx + r$$

$$y' = bx - ny + s$$

$$ay^* - mx^* + r = 0, \quad bx^* - ny^* + s = 0$$

Make Change of Variable $X = x - x^*$, $Y = y - y^*$

Then

$$X' = x' = a(Y + y^*) - m(X + x^*) + r = aY - mX + (ay^* - mx^* + r)$$

$$= aY - mX + 0 = aY - mX$$

Similarly, $Y' = bX - nY$

Write system as

$$X' = aY - mX$$

$$Y' = bX - nY$$

We transform
 $x' = ay - mx + r$
 $y' = bx - ny + s$
a nonhomogeneous system into
 $X' = -mX + aY$
 $Y' = bX - nY$
a homogeneous system.

$$\mathbf{X}' = A \mathbf{X}$$

where

$$A = \begin{bmatrix} -m & a \\ b & -n \end{bmatrix}$$

We can solve by finding eigenvalues and eigenvectors of A

$$\mathbf{X}' = A \mathbf{X}$$

where

$$A = \begin{bmatrix} -m & a \\ b & -n \end{bmatrix} \text{ has solution}$$

$$\alpha e^{\lambda t} \vec{v} + \beta e^{\mu t} \vec{w}$$

where α and β are arbitrary constants

λ is an eigenvalue of A with associated eigenvector \vec{v} and
 $\mu \neq \lambda$ is an eigenvalue of A with associated eigenvector \vec{w} .

The solution of the original system is then

$$\alpha e^{\lambda t} \vec{v} + \beta e^{\mu t} \vec{w} + \begin{bmatrix} x^* \\ y^* \end{bmatrix}$$

Two Particular Examples:

$$\begin{aligned}x' &= -5x + 4y + 1 \\y' &= 3x - 4y + 2\end{aligned}$$

$$(x^*, y^*) = \left(\frac{3}{2}, \frac{13}{8}\right)$$

$$A = \begin{bmatrix} -5 & 4 \\ 3 & -4 \end{bmatrix}$$

$$\lambda = -1, \vec{v} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\mu = -8, \vec{w} = \begin{bmatrix} -4 \\ 3 \end{bmatrix}$$

$$\alpha e^{-t} \begin{bmatrix} 1 \\ 1 \end{bmatrix} + \beta e^{-8t} \begin{bmatrix} -4 \\ 3 \end{bmatrix} + \begin{bmatrix} \frac{3}{2} \\ \frac{13}{8} \end{bmatrix}$$

$$\begin{aligned}x' &= 11y - 9x - 15 \\y' &= 12x - 8y - 60\end{aligned}$$

$$(x^*, y^*) = (13, 12)$$


$$A = \begin{bmatrix} -9 & 11 \\ 12 & -8 \end{bmatrix}$$

$$\lambda = 3, \vec{v} = \begin{bmatrix} 11 \\ 12 \end{bmatrix}$$

$$\mu = -20, \vec{w} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$\alpha e^{3t} \begin{bmatrix} 11 \\ 12 \end{bmatrix} + \beta e^{-20t} \begin{bmatrix} 1 \\ -1 \end{bmatrix} + \begin{bmatrix} 13 \\ 12 \end{bmatrix}$$

$$\begin{aligned}x' &= -5x + 4y + 1 \\y' &= 3x - 4y + 2\end{aligned}$$



$$\begin{aligned}x' &= 11y - 9x - 15 \\y' &= 12x - 8y - 60\end{aligned}$$

